



## Prevalence of Eating Disorders and Disordered Eating Attitudes in Students in Arab Countries: A Scoping Review

TARFA ALBRAHIM\*, ANWAR A. ALRUBAISH, JEWABER T. S. ALFADHLIAH,  
MANAL K. ALASKAR, MONA A. ALATAWI and SARA A. ALDEKHAYIL

Department of Health Sciences, Clinical Nutrition, College of Health and Rehabilitation Sciences,  
Princess Nourah Bint Abdulrahman University, Riyadh 11671, Saudi Arabia.

### Abstract

Anorexia nervosa, bulimia nervosa, and binge-eating disorder represent the specific eating disorders. Although eating disorders (EDs) are a rare psychiatric problem globally, millions of people are exposed to severe health consequences including impairments in health, psychosocial functioning, and quality of life. Young people and adolescents, particularly females, are associated with a higher rate of disordered eating attitudes and EDs than the general population. The aim of this scoping review was to provide an overview or map of ED and disordered eating attitude prevalence in students in Arab countries, both female and male, and to identify contributing factors. Arksey and O'Malley's six-stage framework for scoping reviews was used to guide the process, which returned a total of 20 relevant articles covering 12 of the 22 member countries in the League of Arab States. The average prevalence of ED was 31.4% in a sample of 17,679 students aged 12-33, with a range from 9.6% to 74.5%. Prevalence was higher in female students compared to male students, and the key contributing factors included age, gender, BMI, eating habits, and academic attainment. Given the limitations of this scoping review, especially the lack of a formal critical appraisal process for the included articles, future systematic reviews can use it as a baseline, while empirical researchers and policymakers might pursue its preliminary conclusions.



### Article History

Received: 18 April 2022

Accepted: 23 August 2022

### Keywords

Arab;  
Eating disorders;  
Prevalence;  
Students.

### Introduction

As a group of mental illnesses that affects millions of people worldwide, particularly in Western cultures and young adult and adolescent populations,

eating disorders (EDs) are characterised by complex and damaging relationships with food, and eating.<sup>1</sup> EDs are also associated with distorted body image perceptions, unsuitable weight control

**CONTACT** Tarfa Albrahim ✉ tarfa.ibrahim21@gmail.com 📍 Department of Health Sciences, Clinical Nutrition, College of Health and Rehabilitation Sciences, Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia.



© 2022 The Author(s). Published by Enviro Research Publishers.

This is an Open Access article licensed under a Creative Commons license: Attribution 4.0 International (CC-BY).

Doi: <http://dx.doi.org/10.12944/CRNFSJ.10.2.16>

techniques, and severe purging or eating behaviours (or a combination of both), which lead to insufficient or excessive food intake.<sup>2-3</sup> The mortality rate of EDs is high.<sup>4</sup>

According to the most recent version of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5) eating disorders listed under the category of "Feeding & Eating Disorders" and describes that they are "characterized by a persistent disturbance of eating or eating-related behaviour that results in the altered consumption or absorption of food that significantly impairs physical health or psychosocial functioning."<sup>5</sup> Several categories of the psychiatric condition exist, including binge eating disorder (BED), anorexia nervosa (AN), and bulimia nervosa (BN). Each of these conditions is associated with a unique symptom profile and body weight or shape.<sup>6-7</sup> Specific treatment strategies and long-term, often life-threatening health consequences are also linked to these conditions.<sup>3,8</sup> Disordered eating attitudes, which can be defined as abnormal feelings, behaviours, and beliefs regarding food, are often considered milder types of ED that may or may not warrant a diagnosis of a specific ED. However, disordered eating is significant because they can precede ED, give rise to ED, or indicate ED presence.<sup>9</sup> The term of disordered eating is a descriptive phrase, not a diagnosis. Hence, while many people who have disordered eating may fit the criteria for Eating Disorder Not Otherwise Specified (EDNOS), it also is possible to have disordered eating that do not fit within the current confines of an ED diagnosis.

The lifetime prevalence rates for EDs are relatively low in the general population. However, based on Melisse *et al.*<sup>10</sup> review, 13–55% was at high risk for EDs. On ED-screening instruments, females (11.4–54.8%) displayed more to be at high risk for EDs than males (2–47.3%) with US studies involving female participants reporting that the estimated levels are 1.5% and 0.9% for BN and AN, respectively, which the median age of onset of those disorders among the participants are ranged from 18–21 years.<sup>11</sup> The public health burden associated with these conditions, the severity of their effects on every organ, and the significance of the comorbid conditions associated with EDs have contributed to the growing attention the condition has received in research studies.<sup>12</sup> For example,

health-related outcomes of EDs include poor mental health, digestive disorders, cardiovascular problems, impaired reproductive function, fatigue, substance abuse, gestational diabetes, and pregnancy complications.<sup>8</sup> Furthermore, low bone mineral density (BMD) is a common complication of AN. BMD can develop osteoporosis, or fragile bones, which can cause lifelong debilitating consequences.<sup>13</sup>

An especially critical consideration that has caught the attention of the research community in past years is that, despite the rare nature of EDs in the general population,<sup>1</sup> adolescents, young adults, and college-age individuals, especially females, are at much greater risk of suffering from EDs and disordered eating attitudes. As a case in point, several studies have reported that ED prevalence in college-aged students ranges from 8% to 20.5%, which is significantly higher than the prevalence of these conditions in the general population.<sup>14-17</sup>

An important outcome of the growing level of attention paid in the literature towards EDs and disordered eating attitudes has been the publication of systematic literature reviews (SLRs) seeking to collate and synthesise results from epidemiological studies conducted in specific settings (e.g., world regions or in specific populations such as medical students).<sup>18</sup> Prompted by the recognition that the epidemiology of EDs, as is the case with other health conditions, varies from context to context,<sup>19</sup> SLRs such as Kolar *et al.*<sup>20</sup> and Dahlgren *et al.*<sup>21</sup> have sought to offer accurate estimates of ED prevalence in Latin American and the Nordic countries, respectively. In both cases, the reviewers stressed that accurate information about ED prevalence can play a vital role in evidence-based policy making and practice, particularly regarding the manner in which ED-focused public health interventions should be designed and implemented. Therefore, the aim of this study was to conduct a scoping review that collates, synthesises, and – to the greatest possible extent – critically appraises the results from a growing body of literature pertaining to ED and disordered eating attitude prevalence in students in Arab countries. A cursory literature search indicated that multiple empirical studies have sought to investigate ED prevalence, as well as the risk of disordered eating attitudes, in student populations in Arab countries such as Kuwait, the Kingdom of Saudi Arabia (KSA),

United Arab Emirates (UAE) and Jordan, among others. However, to the best of our knowledge, no previous review study has been conducted to synthesise these findings. Thus, as this scoping review is the first to synthesise results from existing studies relevant to the Arab countries, it has implications for evidence-based policy and further research. Additionally, the scoping review's focus on the factors that contribute to ED and disordered eating attitudes in Arab countries can play a role in guiding healthcare initiatives and future empirical and review studies across the region.<sup>22</sup>

Based on the above, the following research question was established: "What is the prevalence of ED and disordered eating attitudes in students in Arab countries, and what are the contributing factors?". In addition, the aim of the research was decided as follows: "To provide an overview or map of ED and disordered eating attitude prevalence in students in Arab countries, and to identify contributing factors". Finally, to guide the reviewers in addressing this research question and aim, two objectives were established: (i) To synthesise data regarding ED and disordered eating attitude prevalence in students in Arab countries; and (ii) To identify the main factors that can account for ED and disordered eating attitude prevalence in the target population.

## **Methodology**

### **Scoping Review**

Arksey and O'Malley's<sup>23</sup> six-stage framework for scoping reviews was used to guide the process of completing this review. Although refinements have been made to this framework in the years since its publication,<sup>24</sup> and although systematic literature reviews (SLRs) address many of the core limitations associated with scoping reviews, the rationale for choosing a scoping review over an SLR was based on an important consideration. Specifically, the aim was to provide an overview or map of the available evidence regarding ED prevalence in students in Arab countries, given the lack of previous reviews. As Munn *et al.*<sup>25</sup> noted, this is one area in which scoping reviews are valuable, not least because an overview or map of a research issue can inform subsequent systematic reviews.

The six stages of the scoping review framework proposed by Arksey and O'Malley<sup>23</sup> are the following: (i) selecting a research question; (ii)

identifying relevant studies; (iii) selecting relevant studies; (iv) interpreting and synthesising the data; (v) collating, summarising, and reporting the results; and (vi) an optional consultation initiative. With the exception of the final consultation initiative, which typically involves using stakeholder discussions for validation, the rest of the framework was applied. To promote the trustworthiness of the review, which is important given this review's aim of using its results to inform further research and evidence-based policy,<sup>26</sup> the process adopted for this scoping review is described in the rest of this section.

### **Search Strategy**

The electronic databases included in the search were Pub Med, Psych Info, Google Scholar, and MEDLINE. To identify relevant studies for inclusion in this scoping review, the following search term was constructed and entered into the databases without field restriction: ("prevalence") AND ("eating disorder" OR "binge eating disorder" OR "anorexia nervosa" OR "bulimia nervosa" OR "eating disorder not otherwise specified" OR "disordered eating attitudes") AND ("Arab" OR "Gulf Cooperation Council") AND ("young adult" OR "adolescent" OR "university student" OR "student" OR "college student" OR "secondary school").

The search was undertaken in accordance with the PRISMA guidelines, and the selection of relevant studies was informed by the process advocated by Aveyard.<sup>27</sup> Specifically, a reference manager was used to store the search results; duplicate records were excluded; article titles and abstracts were manually screened against the inclusion and exclusion criteria (see Section 2.3); reference lists were searched for further relevant articles; full texts were screened to against the same criteria to assess their eligibility for inclusion; and finally, the remaining articles were included in the scoping review.

### **Inclusion and Exclusion Criteria**

Since search results are likely to contain articles that are unrelated to the research aim,<sup>28</sup> inclusion and exclusion criteria were established in advance to ensure that only relevant articles were considered in the scoping review. To ensure that up-to-date evidence was included in the review, only those articles published after 2012 were included. Additionally, to avoid difficulties arising from the need to translate non-English articles, articles

were included only if they were written in English. Since no formal critical appraisal process was utilised in this scoping review, only peer-reviewed articles were included (quantitative, qualitative, mixed methods, and review studies). Epidemiological measures (e.g. incidence or mortality) and dissertations were excluded.

A key criterion for including articles in this scoping review was that the measurement tool used for assessing ED and disordered eating attitudes in Arabic student populations was a revised version of the eating attitudes test (EAT-26 that including restrictive dieting, fasting, using of laxatives and diuretics self-induced vomiting and/or binge eating). The rationale for this was that EAT-26 has been validated and is considered internally consistent and reliable in both clinical and non-clinical Arabic samples.<sup>29</sup> Additionally, since young people, adolescents, and students are an at-risk population for ED and disordered eating attitudes, only those studies addressing this population were included in the scoping review. Finally, literature was excluded if it did not pertain directly to one of the 22 member countries in the League of Arab States.<sup>30</sup>

## Results

### Search Summary

We have collected 49 studies following the EAT-26 survey. However, 29 papers were excluded due to factors, such as being non-Arab population, age group was younger or older than college age, not using the EAT-26 survey, and being published before 2005. An overview of the studies included in this scoping review ( $n = 20$ ) is given in Table 1. Each study adopted a quantitative, cross-sectional design, which is suitable given the aim of taking a snapshot of the target populations at a particular point in time. Additionally, most were conducted in the KSA ( $n = 10$ ), one study involved 7 Arab countries<sup>31</sup>, and the other countries included in this sample of studies were Kuwait ( $n = 2$ ), Jordan ( $n = 2$ ), Sudan ( $n = 1$ ), Lebanon ( $n = 1$ ), the UAE ( $n = 1$ ), Egypt ( $n = 1$ ), and Morocco ( $n = 1$ ). Therefore, 12 of the 22 member countries in the League of Arab States were addressed in this scoping review.

The total sample size addressed by this scoping review was 17,679 students aged 12-33. The smallest sample ( $n = 100$ ) was associated with Bano *et al.*<sup>32</sup> study and the largest ( $n = 4,698$ ) with

Musaiger *et al.*<sup>31</sup> cross-sectional analysis of students in 7 Arab countries. Most of the studies included both males and females in their sample ( $n = 11$ ), followed by only females ( $n = 8$ ) and only males ( $n = 1$ ). In each case, the outcome measures addressed in the study included ED prevalence or the risk of disordered eating attitudes, and EAT-26, among other measurement tools (e.g., anthropometric measurements of height and BMI, as well as the BSQ and SATAQ-4), were used to collect data.

### Prevalence of ED and Disordered Eating Attitudes

The average prevalence of disordered eating attitudes and ED reported across the 12 included member countries in the League of Arab States was 31.4%. Notably, significant variations in prevalence were observed between the countries. For example, the countries with the highest prevalence were Egypt (73.3%)<sup>33</sup> and Lebanon (74.5%)<sup>34</sup>, but it should be emphasised that only one study was included from each of these settings. The lowest prevalence rate was reported in Morocco (9.6%)<sup>35</sup> but this study only assessed bulimia nervosa (BN), thereby excluding other EDs.

Average prevalence across the studies conducted in the KSA ( $n = 10$ ) amounted to 29.7%, with a range of 21% in Ha'il<sup>32</sup> to 45.5% in Dammam.<sup>36</sup> As for the average prevalence associated with the other countries included in this scoping review, Sudan, Jordan, and the UAE were lower than the KSA at 21.1%, 18.9%, and 20%, respectively. Average prevalence from the studies conducted in Kuwait ( $n = 2$ ) was higher compared to the KSA at 46.4%.

All of the studies which included both females and males reported that the prevalence of ED and disordered eating attitudes were higher in the former population when compared to the latter ( $n = 8$ ). In certain studies, the difference in ED prevalence and disordered eating attitude was significant (e.g., 30% higher for females),<sup>32</sup> while in other studies, the difference was comparatively small (e.g., 5.1% higher for females).<sup>37</sup> For the studies which included only females ( $n = 8$ ), the average prevalence was 30.9%, compared to 20.4% for the only all-male study in this scoping review.

Table 1: Overview of Included Studies

Author(s)	Date	Design	Setting	Population	Sample Size	Age Group (years)	Male/Female /Both	Measurement Tools	Results
Abdelrahim <i>et al.</i>	(2012) <sup>38</sup>	Cross-sectional	Sudan	University students	340	17-23	Both	EAT-26 and Body Shape Questionnaire (BSQ)	21.1% at risk of ED
Bano <i>et al.</i>	(2013) <sup>32</sup>	Cross-sectional	Hail, KSA	Adolescents and young adults	100	18-25	Both	EAT-26	36% and 6% at risk of ED for females and males, respectively
Schulte and Thomas	(2013) <sup>39</sup>	Cross-sectional	UAE	Adolescents	361	19 (mean) age	Both	EAT-26	20% at risk of ED
Musaiger <i>et al.</i>	(2013) <sup>31</sup>	Cross-sectional	Algeria, Jordan, Kuwait, Libya, Palestine, Syria, and UAE	Adolescents	4,698	15-18	Both	EAT-26 and obesity reference	26.7% (Libya), 31.7% (Palestine), 22.9% (Syria), and 33.5% (UAE) at risk of ED
Alihalibi	(2015) <sup>40</sup>	Cross-sectional	Makkah, KSA	Secondary school students	180	15-19	Female	EAT-26	26.1% at risk of ED
Fallatah <i>et al.</i>	(2015) <sup>41</sup>	Cross-sectional	Jeddah, KSA	Secondary school students	425	15-18	Female	EAT-26	32.9% at risk of ED
Kilani	(2017) <sup>37</sup>	Cross-sectional	Jordan	University students	4,565	18-22	Both	EAT-26	14.2% at risk of ED (15.7% and 10.6% for females and males, respectively)

El-Bagoury <i>et al.</i> (2017) <sup>33</sup>	Cross-sectional	Egypt	University students	445	17-26	Both	EAT-26	73.3% at risk of ED
Makdad <i>et al.</i> (2017) <sup>35</sup>	Cross-sectional	Morocco	High school and university students	367	12-19	Both	EAT-26 and hospital scale anxiety and depression (HAD)	9.6% bulimia prevalence (13.5% and 5.3% in females and males, respectively) 45.5% at risk of ED
Alwosaifer <i>et al.</i> (2018) <sup>36</sup>	Cross-sectional	Dammam, KSA	University students	670	18-23	Both	EAT-26 and socio-cultural attitudes towards appearance questionnaire (SATAQ-4) EAT-26	23.6% ED risk (29.4% and 16.4% for females and males, respectively) 35.4% at risk of ED 46.4% at risk of ED
Gharaiabah <i>et al.</i> (2018) <sup>32</sup>	Cross-sectional	Jordan	Secondary school students	738	14-16	Both	EAT-26	30.8% reported binge eating, 7% reported purging, 5.3% reported laxative usage 26.6% at risk of ED 24.1% at risk of ED
Taha <i>et al.</i> (2018) <sup>43</sup>	Cross-sectional	Taif, KSA	University students	1,200	17-33	Female	EAT-26	
Alkazemi <i>et al.</i> (2018) <sup>44</sup>	Cross-sectional	Kuwait	University students	1,147	19-25	Female	EAT-26	
Almuhlaifi <i>et al.</i> (2018) <sup>45</sup>	Cross-sectional	Tabuk, KSA	Secondary school students	480	16 (mean age)	Female	EAT-26	
Fatima <i>et al.</i> (2018b) <sup>46</sup>	Cross-sectional	Arar, KSA	University students	120	18-23	Female	EAT-26	
Fatima <i>et al.</i> (2018a) <sup>47</sup>	Cross-sectional	Arar, KSA	University students	160	18-23	Female	EAT-26	

Fatima and Ahmad	(2018) <sup>48</sup>	Cross-sectional	Arar, KSA	Secondary school students	314	15-19	Female	EAT-26	25.5% at risk of ED
Farchakh <i>et al.</i>	(2019) <sup>34</sup>	Cross-sectional	Lebanon	University students	627	19-24	Both	EAT-26	74.5% at risk of ED
Alhazmi and Al-Johani	(2019) <sup>49</sup>	Cross-sectional	Medina, KSA	University students	342	22 (mean age)	Both	EAT-26	28.7% at risk of ED
Ebrahim <i>et al.</i>	(2019) <sup>50</sup>	Cross-sectional	Kuwait	University students	400	21 (mean)	Male	EA-26	20.4% at risk of ED

## Discussion

After discussing the results from the previous section against those reported elsewhere in the literature, this section draws on evidence from the included studies ( $n = 20$ ) to achieve the second review objective: namely, to identify the main factors that can account for ED and disordered eating attitude prevalence in the target population.

During adolescence and young adulthood, countless studies have demonstrated that individuals are at heightened risk for ED and disordered eating attitudes. This is especially true in females, who often become extremely conscious of their body image, and consequently adopt inappropriate weight loss methods.<sup>1</sup> Empirical studies of students have been published in diverse research contexts, indicating that both females and males are more likely to develop EDs and disordered eating attitudes during adolescence and young adulthood.<sup>21,51</sup> While disordered eating attitudes are not always considered types of ED, they are critical to consider because they represent an important precursor to ED, and they can also indicate the presence of ED.<sup>9</sup>

Therefore, the results of this study regarding the prevalence of ED and the risk of disordered eating attitudes in adolescent and young adult students in Arab countries are largely consistent with the literature. Interestingly, despite the fact that males in Arab countries, including the KSA, are typically associated with greater levels of overweight and obesity,<sup>52</sup> Females were identified as having a higher prevalence of EDs and disordered eating attitudes in the 12 Arab countries included the scoping review. Nevertheless, it is important to emphasise that, similar to the results reported elsewhere in the literature in other research settings, EDs and disordered eating attitudes were still found to affect 5-30% of the participating male students.

The studies included in this scoping review, as well as those published elsewhere in the literature, have highlighted several factors that can account for this study's results regarding the heightened prevalence of ED and disordered eating attitudes in students, particularly in females. Increasing levels of body image consciousness in female students, including those in Arab countries, have been linked

to pressures from mass media and Westernisation, which often imply a connection between beauty standards and thinness.<sup>38,42,45,49-50</sup> In the case of male students, Ebrahim *et al.*<sup>50</sup> emphasised that similar pressures can even lead to body dissatisfaction in this population, but typically for reasons associated with body fat and muscle mass rather than a pre occupation with thinness.

Empirical results from various research settings, including Latin America, Western Europe, and the Nordic countries, have found that strong predictors of ED and disordered eating attitudes in the general population are age and gender.<sup>3,8,18,20-21</sup> The results of this scoping review provide a preliminary confirmation of these results for the Arab countries, particularly given the well-documented, reliable, and valid connections between the female gender, adolescent or young adult student status, and the heightened risk of ED and disordered eating attitudes.

The studies included in this scoping review highlighted several other factors that are correlated with and, to a certain extent, may contribute to ED and disordered eating attitude prevalence. In Alhazmi and Al-Johani<sup>49</sup> KSA-based study, the researchers reported a link between obesity (i.e., BMI) and a high risk of ED, as well as academic attainment and a high risk of ED. While the connection between obesity and ED risk, as an extensively researched issue, has been documented elsewhere in the literature, contrasting results have been reported regarding the connection between academic attainment and ED risk. For example, Alhazmi and Al-Johani<sup>49</sup> found that low academic attainment was associated with an increased risk of ED, while Sundquist *et al.*<sup>53</sup> Sweden-based study reported the opposite. In a sample of Malaysian students, Zuman *et al.*<sup>54</sup> results were consistent with those of Alhazmi and Al-Johani,<sup>49</sup> but further research is certainly warranted in the Arab countries to explore this issue. Finally, three of the included studies documented that, in adolescent females in the KSA, vegetarianism was significantly associated with a greater risk of ED and disordered eating attitudes.<sup>46-48</sup> Evidence suggests that among patients with AN, about 50% report eating some form of vegetarian diet (45-54%), compared to about 6-34%

of adolescent and young adult women overall.<sup>55</sup> The obtained findings confirm those reported in the United States, where vegetarianism in females was identified as a risk factor for ED.<sup>55</sup> While this is an interesting relationship, each of the studies suffers from significant methodological concerns which prevent the identification of confounding variables and the nature of the causal relationship between vegetarianism and EDs. Hence, further exploration of the underexplored factors that may be implicated in EDs and the risk of disordered eating attitudes should be investigated, particularly in the Arab context.

### Conclusion

Eating disorders are potentially life-threatening conditions with severe long-term health consequences. Studies from around the world, including the Arab countries, indicate that ED prevalence is greater in adolescent and young adult populations than it is in the general population. Therefore, understanding the risk factors for EDs, as well as the prevalence of EDs and disordered eating attitudes, is essential, particularly in terms of resource allocation, capacity building, and guiding the design and implementation of evidence-based public health interventions. Although scoping reviews suffer from clear limitations when compared to systematic literature reviews, this review has provided baseline data that future researchers can use as a foundation for further study of this issue in the Arab countries. Conducting an SLR would benefit from a systematic critical appraisal of the studies included in this review and a more comprehensive search strategy. Additionally, future empirical research initiatives could target the key considerations highlighted in this review. In particular, the connection between academic attainment, vegetarianism, and ED risk is an area that merits further attention.

### Acknowledgment

Princess Nourah Bint Abdulrahman University Research Supporting Project number (PNURS2022R69), Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia.



**Funding**

Princess Nourah Bint Abdulrahman University Research Supporting Project number (PNURS2022R69), Princess Nourah Bint Abdulrahman University, Riyadh, Saudi Arabia.

**Conflicts of Interest**

All authors mentioned no conflicts of interest in this research.

**References**

- Blodgett Salafia EH, Jones ME, Haugen EC, Schaefer MK. Perceptions of the causes of eating disorders: a comparison of individuals with and without eating disorders. *J Eat Disord.* 2015/09/15 2015;3(1):32. doi:10.1186/s40337-015-0069-8
- Rikani AA, Choudhry Z, Choudhry AM, *et al.* A critique of the literature on etiology of eating disorders. *Ann Neurosci.* Oct 2013;20(4):157-61. doi:10.5214/ans.0972.7531.200409
- Limbers CA, Cohen LA, Gray BA. Eating disorders in adolescent and young adult males: prevalence, diagnosis, and treatment strategies. *Adolesc Health Med Ther.* 2018;9:111-116. doi:10.2147/AHMT.S147480
- Couturier J, Isserlin L, Norris M, *et al.* Canadian practice guidelines for the treatment of children and adolescents with eating disorders. *J Eat Disord.* 2020;8:4. doi:10.1186/s40337-020-0277-8
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders DSM-5. The American Psychiatric Association (APA) <https://www.psychiatry.org/psychiatrists/practice/dsm>
- Westmoreland P, Krantz MJ, Mehler PS. Medical Complications of Anorexia Nervosa and Bulimia. *Am J Med.* Jan 2016;129(1):30-7. doi:S0002-9343(15)00582-3
- Neyland MKH, Bardone-Cone AM. Treatment experiences of Latinas with current or past binge eating disorder and/or bulimia nervosa. *Eat Disord.* Mar-Apr 2019;27(2):253-265. doi:10.1080/10640266.2019.1591827
- O'Brien KM, Whelan DR, Sandler DP, Hall JE, Weinberg CR. Predictors and long-term health outcomes of eating disorders. *PLoS One.* 2017;12(7):e0181104. doi:10.1371/journal.pone.0181104
- Alvarenga Mdos S, Scagliusi FB, Philippi ST. Development and validity of the Disordered Eating Attitude Scale (DEAS). *Percept Mot Skills.* Apr 2010;110(2):379-95. doi:10.2466/PMS.110.2.379-395
- Melisse B, de Beurs E, van Furth EF. Eating disorders in the Arab world: a literature review. *J Eat Disord.* Nov 6 2020;8(1):59. doi:10.1186/s40337-020-00336-x
- Hudson JI, Hiripi E, Pope HG, Jr., Kessler RC. The prevalence and correlates of eating disorders in the National Comorbidity Survey Replication. *Biol Psychiatry.* Feb 1 2007;61(3):348-58. doi:S0006-3223(06)00474-4
- Kerr-Gaffney J, Harrison A, Tchanturia K. Social anxiety in the eating disorders: a systematic review and meta-analysis. *Psychol Med.* Nov 2018;48(15):2477-2491. doi:S0033291718000752
- Drabkin A, Rothman MS, Wassenaar E, Mascolo M, Mehler PS. Assessment and clinical management of bone disease in adults with eating disorders: a review. *J Eat Disord.* 2017/12/04 2017;5(1):42. doi:10.1186/s40337-017-0172-0
- Eisenberg D, Nicklett EJ, Roeder K, Kirz NE. Eating disorder symptoms among college students: prevalence, persistence, correlates, and treatment-seeking. *J Am Coll Health.* 2011;59(8):700-7. doi:10.1080/07448481.2010.546461
- Tavolacci MP, Grigioni S, Richard L, Meyrignac G, Dechelotte P, Ladner J. Eating Disorders and Associated Health Risks Among University Students. *J Nutr Educ Behav.* Sep-Oct 2015;47(5):412-20 e1. doi:S1499-4046(15)00569-2
- Nivedita N, Sreenivasa G, Sathyanarayana Rao TS, Malini SS. Eating disorders: Prevalence in the student population of

- Mysore, South India. *Indian J Psychiatry*. Oct-Dec 2018;60(4):433-437. doi:10.4103/psychiatry.IndianJPsychiatry\_97\_16
17. Yu Z, Tan M. Disordered Eating Behaviors and Food Addiction among Nutrition Major College Students. *Nutrients*. Oct 26 2016;8(11) doi:nu8110673
  18. Jahrami H, Saif Z, Faris MA, Levine MP. The relationship between risk of eating disorders, age, gender and body mass index in medical students: a meta-regression. *Eat Weight Disord*. Apr 2019;24(2):169-177. doi:10.1007/s40519-018-0618-7
  19. Qian J, Hu Q, Wan Y, *et al.* Prevalence of eating disorders in the general population: a systematic review. *Shanghai Arch Psychiatry*. Aug 2013;25(4):212-23. doi:10.3969/j.issn.1002-0829.2013.04.003
  20. Kolar DR, Rodriguez DL, Chams MM, Hoek HW. Epidemiology of eating disorders in Latin America: a systematic review and meta-analysis. *Curr Opin Psychiatry*. Nov 2016;29(6):363-71. doi:10.1097/YCO.0000000000000279
  21. Lindvall Dahlgren C, Wisting L, Ro O. Feeding and eating disorders in the DSM-5 era: a systematic review of prevalence rates in non-clinical male and female samples. *J Eat Disord*. 2017;5:56. doi:10.1186/s40337-017-0186-7
  22. Azzeh M, Peachey G, Loney T. Prevalence of High-Risk Disordered Eating Amongst Adolescents and Young Adults in the Middle East: A Scoping Review. *Int J Environ Res Public Health*. 2022;19(9)doi:10.3390/ijerph19095234
  23. Arksey H, O'Malley L. Scoping studies: towards a methodological framework. *Int J Soc Res Methodol*. 2005/02/01 2005;8(1):19-32. doi:10.1080/1364557032000119616
  24. Levac D, Colquhoun H, O'Brien KK. Scoping studies: advancing the methodology. *Implement Sci*. Sep 20 2010;5:69. doi:1748-5908-5-69
  25. Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol*. 2018/11/19 2018;18(1):143. doi:10.1186/s12874-018-0611-x
  26. Cairney P, Oliver K. Evidence-based policymaking is not like evidence-based medicine, so how far should you go to bridge the divide between evidence and policy? *Health Res Policy Syst*. 2017/04/26 2017;15(1):35. doi:10.1186/s12961-017-0192-x
  27. Aveyard H. Doing a literature review in health and social care : a practical guide. 2nd ed ed. McGraw-Hill/Open University Press; 2010.
  28. Aveyard H, Sharp P. A Beginners Guide to Evidence Based Practice in Health and Social Care. Maidenhead, Open University Press.; 2009.
  29. Khaled SM, Kimmel L, Le Trung K. Assessing the factor structure and measurement invariance of the eating attitude test (EAT-26) across language and BMI in young Arab women. *J Eat Disord*. 2018;6:14. doi:10.1186/s40337-018-0199-x
  30. Blair I, Grivna M, Sharif AA. The "Arab World" is Not a Useful Concept When Addressing Challenges to Public Health, Public Health Education, and Research in the Middle East. *Front Public Health*. 2014;2:30. doi:10.3389/fpubh.2014.00030
  31. Musaiger AO, Al-Mannai M, Tayyem R, *et al.* Risk of disordered eating attitudes among adolescents in seven Arab countries by gender and obesity: a cross-cultural study. *Appetite*. Jan 2013;60(1):162-167. doi:S0195-6663(12)00426-6
  32. Bano R, Banu D, Shahida S. A Study on the Prevalence and Severity of Eating Disorders among the Young Population of Ha'il City in Saudi Arabia. *Glob J Res Anal*. 2013;2(6)
  33. El-Bagoury L, Hassan A, AbouSeif H. Eating attitudes and barriers to healthy eating and physical activity among a sample of university students in Egypt. *J Egypt Public Health Assoc*. 2017;92 (1):29-35.
  34. Farchakh Y, Hallit S, Soufia M. Association between orthorexia nervosa, eating attitudes and anxiety among medical students in Lebanese universities: results of a cross-sectional study. *Eat Weight Disord*. Aug 2019;24(4):683-691. doi:10.1007/s40519-

- 019-00724-6
35. Makdad M, Najdi A, Bour A. Risk Factors Associated with Bulimia in Schooled Adolescents in Fez (Morocco). *Eur J Sci Res.* 2017;148 (1):155-160.
  36. Alwosaifer AM, Alawadh SA, Abdel Wahab MM, Boubshait LA, Almutairi BA. Eating disorders and associated risk factors among Imam Abdulrahman bin Faisal university preparatory year female students in Kingdom of Saudi Arabia. *Saudi Med J.* Sep 2018;39(9):910-921. doi:10.15537/smj.2018.9.23314
  37. Kilani BF. Prevalence of Eating Disorders among Students at the University of Jordan. *Arab J Psychiatry.* 2017;28 (1):76-82.
  38. Abdelrahim F, Abdelmutti N, Alshaikh A. Eating disorders symptoms among a university students: an exploratory study. *Sudan Med J.* 2012;48 (3):186-193.
  39. Schulte SJ, Thomas J. Relationship between eating pathology, body dissatisfaction and depressive symptoms among male and female adolescents in the United Arab Emirates. *Eat Behav.* 2013/04/01/2013;14(2):157-160. doi:https://doi.org/10.1016/j.eatbeh.2013.01.015
  40. Allihaibi MM. Disordered eating attitudes among secondary schoolgirls in Al-Iskan sector, Makkah Al-Mukarramah, Saudi Arabia. *Int J Med Sci Public Health.* 2015;4(7):939-945.
  41. Fallatah A, Al-Hemairy M, Al-Ghamidi H. Eating Disorders among Female Adolescents in Jeddah. Scientific Cooperation Medical Workshops. .2015; Accessed 25 October 2019. [https://pdfs.semanticscholar.org/f16e/714064b38d0650f9dec91551cc8a75ab05f3.pdf?\\_ga=2.195505580.1424072328.1572564845-3276361.1571329572](https://pdfs.semanticscholar.org/f16e/714064b38d0650f9dec91551cc8a75ab05f3.pdf?_ga=2.195505580.1424072328.1572564845-3276361.1571329572)
  42. Gharaibah T, Al-Sheyab N, Shiyab W. Prevalence and Correlates of Disordered Eating Attitudes Among Jordanian Adolescents. *Arab J Nutr Exerc.* 2018;3 (2):35-53.
  43. Taha A, Hany A, Desouky D. Eating Disorders Among Female Students of Taif University, Saudi Arabia. *Arch Iran Med.* 2018;21 (3):111-117.
  44. Alkazemi D, Zafar TA, Ebrahim M, Kubow S. Distorted weight perception correlates with disordered eating attitudes in Kuwaiti college women. *Int J Eat Disord.* May 2018;51(5):449-458. doi:10.1002/eat.22852
  45. Almuhlafi M, Jamilah KA, Almutairi AF, Salam M. Relationship between early menarche, obesity, and disordered eating behaviors: a school-based cross-sectional survey in Northern Saudi Arabia. *Diabetes Metab Syndr Obes.* 2018;11:743-751. doi:10.2147/DMSO.S180697
  46. Fatima W, Fatima R, Anwar N. Subclinical eating disorders and association with vegetarianism in female students of Saudi Arabia. A cross-sectional study. *IOSR J Nurs Health Sci.* 2018;7(2):62-67.
  47. Fatima W, Fatima R, Anwar N. Prevalence of Eating Disorders among Female College Students of Northern Broader University, Arar, Kingdom of Saudi Arabia. *Int J Child Health Nutr.* 2018;7(115-121)
  48. Fatima W, Ahmad L. Prevalence of disordered eating attitudes among adolescent girls in Arar City, Kingdom of Saudi Arabia. *Health Psychol Res.* 2018;6(1):7444.
  49. Alhazmi A, Al-Johani A. Prevalence and associated factors of eating disorders among students in Taiba University, Saudi Arabia: A cross-sectional study. *Malaysian J Public Health Med.* 2019;19 (1):172-176.
  50. Ebrahim M, Alkazemi D, Zafar TA, Kubow S. Disordered eating attitudes correlate with body dissatisfaction among Kuwaiti male college students. *J Eat Disord.* 2019;7:37. doi:10.1186/s40337-019-0265-z
  51. Ganesan S, Ravishankar S, Ramalingam S. Are Body Image Issues Affecting Our Adolescents? A Cross-sectional Study among College Going Adolescent Girls. *Indian J Community Med.* 2018;43 (1):42-46.
  52. Alkazemi D. Gender differences in weight status, dietary habits, and health attitudes among college students in Kuwait: A cross-sectional study. *Nutr Health.* Jun 2019;25(2):75-84. doi:10.1177/0260106018817410
  53. Sundquist J, Ohlsson H, Winkleby MA, Sundquist K, Crump C. School Achievement and Risk of Eating Disorders in a Swedish

- National Cohort. *J Am Acad Child Adolesc Psychiatry*. Jan 2016;55(1):41-46 e1. doi:S0890-8567(15)00652-8
54. Zuman N, Ilias K, Isa K, Danis A. Relationship Between Eating Behaviors, Self Esteem and Academic Achievement among Lower Secondary School Students in Meru Klang, Malaysia. *Asian Journal of Clinical Nutrition*. 2012;4(4):132-141.
55. Bardone-Cone AM, Fitzsimmons-Craft EE, Harney MB, *et al.* The inter-relationships between vegetarianism and eating disorders among females. *J Acad Nutr Diet*. Aug 2012;112(8):1247-52. doi:S2212-2672(12)00627-2