

An Analysis of Contributors to Energy Intake Among Middle Aged and Elderly Adults

NATASHA HURREE¹ and RAJESH JEEWON^{1*}

Department of Health Sciences, Faculty of Science, University of Mauritius, Réduit, Mauritius.

<http://dx.doi.org/10.12944/CRNFSJ.4.Special-Issue-Elderly-November.02>

(Received: June 29, 2016; Accepted: July 15, 2016)

ABSTRACT

Adulthood and middle age is widely recognized as the time of life when unhealthy eating habits may develop. Data from various studies have demonstrated that changes in eating habits may also occur during old age. It is essential to acknowledge that a high consumption of certain food groups such as sweetened beverages, meat and eggs may contribute to an increased energy intake. This obviously results in high body mass index (BMI) and consequently an increased risk of non-communicable diseases (NCDs) and obesity. Energy intake among middle aged and elderly individuals may be influenced by socio demographic factors (for example: age, gender, socio economic status), social factors (for example: marital status), environmental factors like access to food commodities as well as nutrition knowledge and physical activity level. The present review highlights eating habits, contribution of specific food groups to energy intake and the influence of several factors on energy intake among the middle aged and elderly population.

Keywords: Obesity, eating habits, socio economic status, middle age, elderly.

INTRODUCTION

General eating habits

It is well documented that middle age is the most productive period of life in terms of work capacity but is also the time when unhealthy eating habits and sedentary lifestyles may develop, resulting in malnutrition and increased risk of chronic diseases¹. Intake of fat, animal products and sugar is increasing while on the other hand, consumption of cereals, fruits and vegetables is decreasing across developing countries². Similar results have been found among middle aged adults where a shift from cereals, pulses and potatoes to meat, fats, sugar, cheeses and desserts was noted³. Eating habits tend to vary between gender and across countries as well. For instance, middle aged individuals are found to consume rice, fish and vegetables on a nearly daily basis in Malaysia, with men consuming more vegetables than women⁴. However, in France, middle aged women have a higher consumption of fruits and

vegetables than men⁵. Men tend to consume red and processed meat more often than women while the latter consume skimmed milk and high fiber cereals in greater amounts⁶. It has been reported that food habits of middle aged individuals tend to differ from those of elderly people and that the latter skipped meals less often than middle aged adults⁴. With ageing, people tend to reduce the consumption of meat and certain dairy products but the intake of fish is likely to increase⁷.

Old age is accompanied by several transitions such as retirement which may impact on nutrition^{8,9}. Physiological changes which reduce hunger in the elderly may result in decreased energy intake¹⁰. This finding is supported from a study in USA where older adults were found to have low energy intakes¹¹. Generally, older adults tend to be less active and hence, there is a need to diminish energy intake due to a decrease in Basal Metabolic Rate (BMR)¹² but some believe they do not need

to adjust the quantity of food intake as they are already eating properly¹³. This can result in obesity as reported in the US where the percentage of obesity among the elderly has increased from 14.7 in 1991 to 22.9% in 2000¹⁴. A rise in obesity from 13 to 27% in elderly people has also been noted in China¹⁵.

During old age weight tends to decrease¹⁶ while during middle age body weight increases¹⁷. The main reason is due to an energy imbalance whereby energy intake exceeds energy expenditure¹⁸. The energy requirements of middle aged men of 30-60 years who do moderate physical activity is 2761 kcal/ day while those of women in the same age range is 2225 kcal/day. Energy requirements are generally higher for middle aged than elderly people¹⁹. It has also been postulated that the reduction of energy dense foods combined with increases in fruit and seafood intakes can lead to a decline in Body Mass Index (BMI)²⁰. This review paper aims to elaborate on some of the factors which can influence energy intake among adults and elderly individuals.

Contribution of food groups to energy intake

Recently, the emphasis has been on the study of different food groups which exist and their relationships with energy intake and hence, on the prevalence of overweight and obesity. Investigations are no longer restricted on individual nutrients or food groups since it has been recognized that food items are nearly always consumed in combination. The different food groups can be classified into: grains, proteins, dairy, fruits, vegetables and oils²¹.

In Teheran, a shift towards the western dietary pattern which includes the frequent consumption of processed foods, sweets and fatty food items is linked to an increased energy intake and hence, a high BMI²². It was also observed that a mixed dietary pattern which includes sweets, solid fats and meats as well as healthy items such as nuts, seeds and legumes does not lead to an increased BMI due to the possible interaction of the healthy components to compensate for the effect on obesity²².

Several studies reported the effects of different food groups on BMI and energy intake. The 1999-2004 National Health And Nutrition

Examination Survey (NHANES) during 1999 to 2004 demonstrated that a high consumption of whole grains such as rice, oats, and brown rice was inversely correlated with BMI and the prevalence of obesity²³. Similar results were obtained in Canada where a high carbohydrate intake was linked to low energy intake²⁴. The consumption of barley was found to enhance satiety and decrease energy intake more than whole grain wheat and refined rice in adults and especially among the elderly²⁵. However, it has been suggested that the inverse relationship between high carbohydrate intake and low energy intake and BMI may not be related to lower total energy intake²⁶ but may be due to higher physical activity level²⁷.

There have been some controversial findings as to whether non-vegetarian or vegetarian diet is linked to BMI. Meats, considered as being high in energy and fat may be linked to a higher prevalence of obesity²⁸. Support for this claim comes from findings which indicate that high intakes of meat are associated with a high BMI and Waist Circumference (WC). In the same context, 'meat fat' (red meat, oil, poultry and processed meats) dietary pattern has a positive correlation with obesity in middle aged and elderly people whereas 'vegetables-seafood' (fish, soy foods, vegetables, fruits and sea weeds) dietary pattern are not linked to a high BMI and obesity²⁹. Similar results are reported in the Adventist study where BMI gain are not found to be linked to vegetarian diets as compared to a non-vegetarian diet³⁰. Following a vegetarian diet results in lower energy intake, higher fiber content and lower animal fats intake than non-vegetarian diet^{30, 31}. BMI values are higher in non-vegetarians than their vegetarian counterparts in all age groups and for both men and women³².

On the other hand, no significant differences were reported in energy intake between vegetarians and non-vegetarians and the BMI of male vegetarians did not differ significantly from that of their non-vegetarian counterparts³³. There is no evidence to suggest that one protein source is preferable than another but it has been advocated that animal proteins, especially dairy can support muscle protein synthesis better than plant proteins which can enhance energy expenditure³⁴.

In the US, energy beverages such as soft drinks are on the rise and are replacing water and milk. Energy beverages largely contribute to high energy intake which then leads to weight gain³⁵. In the US, one in four people get at least 200 calories from sugary drinks and individuals who increase their intake of sugary beverages by one 12-ounce serving per day gain much more weight than those who do not alter their consumption³⁶. However, further research between sugar sweetened beverages and weight gain is required³⁷. Alcohol is another beverage from which individuals can derive energy; in the U.S, the adult population consumes an average of 100 calories per day from alcoholic drinks, with the highest prevalence among men than women and younger adults aged 20-39 years than older ones³⁸. About 527 kcal/day of energy was obtained from alcohol among Mexican adults³⁹. High alcohol consumption is associated with poor dietary habits⁴⁰. However, epidemiologic evidence does not show a clear cut relationship between alcohol intake and bodyweight and this complex association needs to be further studied⁴¹.

Major factors influencing energy intake

Besides the normal process of ageing and changes which accompany it such as sarcopenia, there are diverse factors which can influence energy intake of individuals⁴².

Gender

The biochemistry and physiology of women and men differ⁴³. It has been reported that women, generally have lower energy intakes than men⁴⁴ due to the fact that women have a lower body mass and fat free mass but a higher % of body fat⁴⁵. Females tend to choose more nutritious food items than males and have been found to be more concerned about health issues^{46, 47}. De Silva *et al.* (2011) also attributed the gender differences in food consumption to stronger beliefs of women concerning healthy eating⁴⁸. Moreover, women have better awareness and knowledge which may further contribute to healthy eating⁴⁹. An alternative explanation may be that females are more likely to engage in weight control behaviors⁵⁰ and are more willing to compromise on taste to meet their weight related goals⁵¹. In addition, certain foods like red meat, alcohol and large portion sizes have been

associated with masculinity while vegetables, fruit and dairy products are linked to femininity⁴³. However, some men do express some interest in nutrition and health and also advocate that they are trying to reduce red meat consumption and simultaneously, adding more vegetables to their diets⁴³.

Socio economic status (SES)/ Income level

It is well known that socio economic factors have a strong influence on the diet and energy intake⁵². In general, people from higher socioeconomic groups are found to have a healthier dietary intake while poor people are more likely to become obese due to less healthy eating habits⁵³. A study⁵⁴ in Mauritius also reported that the diet quality of women from medium and high SES had better diet quality than those of lower SES. The high energy intake of low income people has been attributed to the low cost of energy dense foods^{17, 55}. Low-cost energy dense foods are both energy rich and shelf stable as compared to low energy dense diets which cost more⁵⁶. Although it has been reported that low income people tend to become overweight, this may not always be true. For instance, chronic energy deficiency was found to be significantly higher among individuals of low socioeconomic status (SES) than those in high SES⁵⁷. Across Mexico, a negative association between SES and obesity has been reported². It has been suggested that an inverted U-shape relationship may exist between socio economic status and obesity as those in extreme poverty do not have resources to become obese while individuals who are slightly well off but still poorer as compared to the population, could maintain a positive energy balance for a long span of time². People from disadvantaged socio economic background tend to practice more unhealthy behaviors due to low self-esteem which lead to greater health complications^{58, 59}.

Contradictory results have been reported whereby low SES individuals have been found to choose nutritious foods⁵⁸. In Canada, higher income individuals are more likely to follow dietary recommendations due to greater awareness of relationships between diet and diseases and are also less resistant to change⁶⁰. Obesity has also been found to be more prevalent in higher socio economic groups in rural India⁶¹.

In developed countries, SES was inversely related to obesity in women but not in men⁶². Among Chinese men, a positive relation was identified between income and obesity and in developing countries, a positive correlation between obesity and SES was noted in both gender⁶². Fast food consumption was also reported to be higher in low income individuals since they have easier access to fast food restaurants than to supermarkets where fresh foods are available, resulting in high frequency of fast food consumption⁶³. It is noted, however, that both high and low income individuals rely on fast foods and hence, have higher energy intakes⁶⁴.

Snacking

Meal and snack patterns have been linked with energy intake among US adults⁶⁵. Total snacking was found to increase about 24% in all adults, with a higher increase observed from 1997-1978 to 2003-2006⁶⁶. Moreover, sweet and fatty food groups are linked with snacking⁶⁷. While younger Malaysian adults derive high energy intakes from morning meals and dinner, older adults have lower intakes from these meals⁶⁸. Among older adults, the prevalence of snacking is 84% and contributes a quarter of their daily energy, with each snacking occasion contributing approximately 150 kcal¹¹. Duffey *et al.* (2013) also reported that in Brazil, snacking is a major source of daily energy intake with the lowest % of daily energy intake from snacks among the 40-59 age group⁶⁹. Multiple snackers have higher energy intakes than those who never snack⁷⁰. However, the notion that high energy intake from snacks contributes to weight gain has been challenged by Drummond *et al.* (1996) and Dongen *et al.* (2010) who advocated that snacking may not necessarily result in weight gain^{71, 72, 73}. One plausible reason is because people can opt for snacks containing high fiber foods such as dried prunes that tend to increase satiety and decrease energy intake⁷⁴.

Portion sizes and access to foods

Food service portion sizes, including fast food and ready-to-eat foods have increased substantially over the past 20 years and it is most likely that individuals are unaware of the amount of food they consume⁷⁵. Portion sizes are common everywhere from restaurants to supermarkets to vending machines⁷⁶ and consumers eat 30 to 50 %

more from larger-sized restaurant portions compared to smaller portions⁷⁵. When portion size and energy density are increased, they act synergistically to increase energy intake⁷⁶. Chronic exposure to large portion sizes was found to result in a rise in energy intake and consequently, weight gain over time among middle aged individuals⁷⁷. A 50% increase in portion size can cause a rise in daily energy intake by 16% in middle aged adults⁷⁸. Moreover, a high prevalence of obesity has been reported with quick service restaurants⁷⁶. Increases in body weight and fatness are also associated with high consumption of fast foods which leads to increased energy intakes in middle aged women^{63, 79}. However, so far this trend has been restricted to the US and western countries and it is unclear whether the frequency of eating out is related to high BMI in Asian countries as well⁸⁰. It can also be argued that the effect of portion size on energy intake has little effect on body weight since individuals may compensate for their high food intake by eating less at the following meal⁸¹.

Food accessibility may be a factor contributing to greater consumption of food and hence, a high energy intake. The more accessible a food, the more it is consumed⁷⁵. In some rural areas, access to transport is difficult and food insecurity may result due to poor health and limited food availability owing to inadequate grocery stores in the neighborhood. Physical disabilities may also impede food intake related activities, such as shopping and cooking, leading to pronounced consequences on eating behavior⁸² but Dressler *et al.* (2013) reported that food security and BMI are not necessarily related⁸³.

Social factors

In developed countries, a positive association between marriage and BMI was demonstrated⁸⁴. The same trend was observed in developing countries as well⁸⁴. In Jamaica, both men and women who were married were found to have higher BMI. This has been attributed to the fact that important events in life may lead to positive energy balance and precipitate the onset of obesity⁸⁵. Among Chinese adults, marital status was positively related to BMI in men⁵². While the dietary food habits of people who were never married consisted of potatoes and red meat, those who were married ate more fish, nuts and legumes. Those who were never married also consumed

more fast foods and drank less alcohol than married or widowed individuals; however, no significant relationship was reported between marital status, BMI and cholesterol levels⁸⁶. Similarly, with reference to marital status, no differences were observed for both males and females in terms of obesity⁸⁷.

While living alone had a great negative impact on dietary habits and nutrient intakes of men, women on the other hand, were found to have nutrient intakes equal or greater than those of women living with a spouse⁸⁸. People living alone were not only at higher nutritional risk but also consumed less meals, had lower intakes of protein, fruits and vegetables as well as a low BMI¹⁰. Moreover, elderly individuals who live alone are more at risk for anorexia of aging and other health problems¹⁰. Widowhood results in higher vulnerability for under nutrition due to reduced food intake as a result of decreased participation in social activities, changes in eating patterns and in social relationships¹⁰. Widowhood may also result in social isolation which can reduce social contact and increase risk for the development of depression. Social isolation results in a higher number of meals eaten alone, which was found to be associated with 30% lower energy intake¹⁰.

Physical Activity Level

Physical activity, considered as an important determinant of body weight can prevent obesity⁸⁹. PA is known to contribute to an improvement in energy and macronutrient balance regulation⁹⁰.

Globally, more than 30% of adults are not sufficiently active⁹¹. Physical activity is closely interrelated with energy intake⁹² as it has the ability to manipulate energy expenditure and when performed on a regular basis, it leads either to loss of body weight or to a need for an increase in energy intake⁹². Weight loss may not occur if people increase their caloric intake after exercise but the extent to which this compensation takes place remains unresolved⁹³. An increase in physical activity leads to a rise in food intake so that energy balance is maintained, but the contrary is not true⁹⁴. It has also been observed that people engaged in physical activity regularly are less likely to gain weight. Among Malay adults a negative association between moderate to high physical activity level and metabolic syndrome was reported⁹⁵. In Mauritius, high physical activity

level is linked to better eating habits among older adults but middle aged women who were engaged in moderate physical activity were found to be overweight⁹. Physical activity is known to decrease with age⁹⁶. However, Indian women aged 51-60 were found to engage in heavy activities as compared to the other age groups (middle aged and young women)⁹⁷. Overweight individuals were reported to have lower physical activity than their normal weight counterparts and small significant relationships were obtained between body fatness and physical activity, however, more research is required to establish a strong link⁹⁸. Moreover, significant associations between sedentary behaviors and weight gain are small⁹⁹ and Donnelly *et al.* (2014) argued that physical activity level does not have any effect on energy intake¹⁰⁰. Physical activity may also differ according to gender as demonstrated in Korean Americans, where men are more engaged in frequent physical activity than women in late life¹⁰¹. In another study, physical activity was negatively associated to BMI in women but not in men⁵⁸.

Nutrition Knowledge (NK)

Education may be associated with increased NK as well as with the ability to translate NK into better health practices¹⁰². In a study, it was observed that a high proportion of older adults in England had little basic NK which was a barrier to healthier eating¹³. People with a good NK are likely to make healthier food choices. For instance, elders' NK of dairy products predicted the type of milk consumed¹³. However, nutritional knowledge is not always consistent with healthy eating habits or adequate energy intake as shown in a study where consumption of several food groups increased with high nutrition knowledge except for fruits and vegetables⁹. A high NK can enable people to improve their nutrition practices and establish desirable eating habits to maintain health¹⁰³. Women have higher NK scores and more negative views of fatty foods, sweets and salty food items than men¹³. Knowledge of fat nutrition in middle aged women led to consumption of low fat diets⁴⁷. Participants having lower education and hence, most probably lower NK consumed less fish and vegetables but more fried foods, pasta and potatoes¹⁰⁴. However, no negative association was reported between nutrition knowledge and fat intakes¹⁰⁵. High nutrition knowledge may not necessarily lead to a healthy

diet [106]. Where obese people are concerned, they may find it difficult to translate knowledge into practice¹⁰⁷.

Dental health/oral status

Dentition has been found to form part of the factors that can increase the risk of malnutrition in elderly people¹⁰⁸. A similar observation was made in another study where tooth loss mouth hygiene, xerostomia and periodontal disease were found to be related to a decrease in taste sensitivity¹⁰. A decrease in sensitivity usually results in decreased food consumption and eventually to the anorexia of aging^{10, 109}. Elderly people may be subject to dietary restrictions due to oral problems such as fewer natural teeth, impaired dentition or dental caries¹¹⁰. Elderly people having compromised functional dentition were reported to have inadequate calorie intake^{111, 113}. While a significant association was reported between dental status and dietary intakes in elderly individuals¹¹², this was found to be more factual for micronutrients but true to a lesser extent for energy intake¹¹². Moreover, despite having dental

impairment or chewing problems, elderly people may not necessarily exclude any food type from their diets¹¹⁴.

CONCLUSION

This review provides insights into the various factors which may impact on energy intake of middle aged and elderly individuals and there is ample evidence to suggest that the factors discussed do have an influence on energy intake. It is also evident from published studies that eating habits with regular consumption of certain food groups have an effect on energy intake. Although evidence does point out that these factors are associated with energy intake, further research and follow up studies are necessary to establish stronger associations between energy intake and the factors discussed. It is also useful to consider these factors while devising new strategies or nutrition interventions to address issues on energy intake, eating habits and obesity among the middle aged and elderly population.

REFERENCES

1. Driskell J.A, *Nutrition and Exercise Concerns of Middle Age*. United States of America: CRC Press (2009).
2. Fernald L.C.H. Socio-economic status and body mass index in low-income Mexican adults. *Social Science and Medicine*; **64**: 2030-2042: (2007).
3. Tyrovolas S, Polychronopoulos E. Lessons from Studies in Middle-Aged and Older Adults Living in Mediterranean Islands: The Role of Dietary Habits and Nutrition Services. *Cardiology Research and Practice*; 1-7: (2011).
4. Karim N.A, Kather H.M.M. Nutritional Status and Food Habits of Middle-aged Adults in Selected Areas of Selangor. *Malaysian Journal of Nutrition*; **9**(2): 125-136: (2003).
5. Estaquio C, Druesne-Pecollo N, Latino-Martel P et al. Socioeconomic Differences in Fruit and Vegetable Consumption among Middle-Aged French Adults: Adherence to the 5 A Day Recommendation. *Journal of the American Dietetic Association*; **108**: 2021-2030: (2008).
6. Fleming S, Kelleher C, O'Connor M. Eating patterns and factors influencing likely change in the workplace in Ireland. *Health Promotion International*; **12**(3):187-196: (1997).
7. Wahlqvist ML and Lukito W, *Nutrition in the Elderly*. Australia: ExcerptaMedica Pty Limited (1992).
8. MC Naughton S.A, Crawford D, Ball K et al. Understanding determinants of nutrition, physical activity and quality of life among older adults: the Wellbeing, Eating and Exercise for a Long Life (WELL) study. *Health and Quality of Life Outcomes*; **109**(10): 1-7: (2012).
9. Dunneram Y, Ramasawmy D, Pugo-gunsam P et al. Determinants of eating habits among pre-retired and post-retired Mauritians. *International Journal of Food Sciences and Nutrition*; **2**(3): 109-115:(2013).
10. De Boer A, Horst G.J.T, Lorist M.M.

- Physiological and psychosocial age-related changes associated with reduced food intake in older persons. *Ageing Research Reviews*; **12**: 316-328: (2013).
11. Zizza C.A, Tayie F.A and Lino M. Benefits of Snacking in Older Americans. *Journal of the American Dietetic Association*; **107**: 800-806 : (2007).
 12. Stanga Z. Basics in clinical nutrition: Nutrition in the elderly. *European e-journal of clinical nutrition and metabolism*; **4**: 289-299: (2009).
 13. Al Riyami A, Al Hadabi S, Abdelaty M.A et al. Nutrition knowledge, beliefs and dietary habits among elderly people in Nizwa, Oman: implications for policy. *Eastern Mediterranean Health Journal*; **16**(8): 859-867: (2010).
 14. Villareal D.T, Apovian C.M, Kushner R.F et al. Obesity in older adults: technical review and position statement of the American Society for Nutrition and NAASO, The Obesity Society. *American Journal of Clinical Nutrition*; **82**: 923-934: (2005).
 15. Chapman I.M. Obesity Paradox during Aging. *Interdisciplinary Topics In Gerontology*; **37**: 20-36: (2010).
 16. Apostolopoulou M, Savopoulos C, Michalakis K et al. Age, weight and obesity. *Maturitas*; **71**: 115-119: (2012).
 17. Baum C.L, Ruhm C.J. Age, socioeconomic status and obesity growth. *Journal of Health Economics*; **28**: 635-648:(2009).
 18. Gill T. Importance of preventing weight gain in adulthood. *Asia Pacific Journal of Clinical Nutrition*; **11**(Suppl): 632-636: (2002).
 19. National Institute Of Nutrition (2009) *Nutrient requirements and recommended dietary allowances for Indians*. Hyderabad, India: Indian Council of Medical Research.
 20. Nakade M, Lee J.S, Kawakubo K et al. Correlation between food intake change patterns and body weight loss in middle-aged women in Japan. *Obesity Research and Clinical Practice*; **1**: 79-89: (2007).
 21. United States Department Of Agriculture (2013) Food groups. Available at: <http://www.choosemyplate.gov/food-groups/protein-foods.html> (accessed 15 October 2013).
 22. Hosseini-Esfahani F, Djazaieri S, Mirmiran P et al. Which Food Patterns Are Predictors of Obesity in Tehranian Adults?. *Journal of Nutrition Education and Behavior*; **44** (6): 564-573: (2012).
 23. O'Neil C.E, Zanutta M, Cho S.S et al. Whole grain and fiber consumption are associated with lower body weight measures in US adults: National Health and Nutrition Examination Survey 1999-2004. *Nutrition Research*; **30**: 815-822: (2010).
 24. Merchant A.T, Vatanparast H, Barlas S et al. Carbohydrate Intake and Overweight and Obesity among Healthy Adults. *Journal of the American Dietetic Association*; **109**: 1165-1172: (2009).
 25. Schroeder N, Gallaher D.D, Arndt E.A et al. Influence of whole grain barley, whole grain wheat, and refined rice-based foods on short-term satiety and energy intake. *Appetite*; **53**: 363-369: (2009).
 26. Wu S, Chang Y, Wei I et al. Intake levels and major food sources of energy and nutrients in the Taiwanese elderly. *Asia Pacific Journal of Clinical Nutrition*; **14**(3): 211-220: (2005).
 27. Gaesser G.A. Carbohydrate Quantity and Quality in Relation to Body Mass Index. *Journal of the American Dietetic Association*; **107**: 1768-1780: (2007).
 28. Wang Y, Beydoun M.A. Meat consumption is associated with obesity and central obesity among US adults. *International Journal of Obesity*; **33**(6): 621-628: (2009).
 29. Cho Y.A.E, Shin A, Kim J. Dietary Patterns Are Associated with Body Mass Index in a Korean Population. *Journal of the American Dietetic Association*; **111**: 1182-1186: (2011).
 30. Japas C, Knutsen S, Dehomb S et al. Body mass index gain between ages 20and 40 years and lifestyle characteristics of men at ages 40—60 years: The Adventist Health Study-2. *Obesity Research and Clinical Practice*; **1**-9: (2013).
 31. Rosell M, Appleby P, Spencer E et al. Weight gain over 5 years in 21 966 meat-eating, fish-eating, vegetarian, and vegan men and women in EPIC-Oxford. *International Journal of Obesity*; **30**:1389–1396: (2006).
 32. American Dietetic Association. Position of the American Dietetic Association: Vegetarian Diets. *Journal of the American Dietetic Association*; **109**: 1266-1282: (2009).

- 15 HURREE & JEEWON, *Curr. Res. Nutr Food Sci Jour.*, Vol. 4(Special Issue 3), 08-18 (2016)
33. Bedford J.L, Barr S.I. Diets and selected lifestyle practices of self-defined adult vegetarians from a population-based sample suggest they are more 'health conscious'. *International Journal of Behavioral Nutrition and Physical Activity*; **4**(2): 1-11: (2005).
34. Gilbert J.A, Bendsen N.T, Tremblay A et al. Effect of proteins from different sources on body composition. *Nutrition, Metabolism and Cardiovascular Diseases*; **21**: 16-31: (2011).
35. Dennis E.A, Flack K.D, Davy B.M. Beverage consumption and adult weight management: A review. *Eating behaviours*; **10**: 237-246: (2009).
36. Harvard School Of Public Health (2012) Fact sheet: sugary drink supersizing and the obesity epidemic. *United States of America: Department of Nutrition at Harvard School of Public Health 2012*.
37. Malik V.S, Schulze M.B and Hu F.B. Intake of sugar-sweetened beverages and weight gain: a systematic review. *American Journal of Clinical Nutrition*; **84**: 274-288: (2006).
38. Nielsen S.J, Siega-Riz A.M and Popkin B.M. Trends in Energy Intake in U.S. between 1977 and 1996: Similar Shifts Seen across Age Groups. *Obesity Research*; **10**(5): 370-378: (2002).
39. Barquera S, Hernandez-Barrera L, Tolentino ML et al. Energy Intake from Beverages Is Increasing among Mexican Adolescents and Adults. *Nutrition Journal*; **138**: 2454-2461: (2008).
40. Herbeth B, Samara A, Stathopoulou M et al. Alcohol Consumption, Beverage Preference, and Diet in Middle-Aged Men from the STANISLAS Study. *Journal of Nutrition and Metabolism*; 1-6: (2012).
41. Jéquier E. Alcohol intake and body weight: a paradox. *American Journal of Clinical Nutrition*; **69**: 173-174: (1999).
42. Ministry Of Health, New Zealand (2013) Food and Nutrition Guidelines for Healthy Older People: A background paper. *Wellington, New Zealand: Ministry of Health*.
43. Arganini C, Saba A, Comitato R et al. Gender Differences in Food Choice and Dietary Intake in Modern Western Societies. In: Maddock J (eds) *Public Health - Social and Behavioral Health Europe*, pp. 83-102: (2012).
44. Kusumaratna R.K. Gender differences in nutritional intake and status in healthy free-living elderly. *UniversaMedicina*; **27**(3): (2008).
45. Tarnopolsky M (1998) *Gender Differences in Metabolism: Practical and Nutritional Implications*. United States of America: CRC Press LLC.
46. Beardsworth A, Bryman A, Keil T, Goode J et al. Women, men and food: the significance of gender for nutritional attitudes and choices. *British Food Journal*; **104**(7): 470 – 491:(2002).
47. Worsley A, Wang W.C, Hunter W. Gender differences in the influence of food safety and health concerns on dietary and physical activity habits. *Food Policy*; **41**: 184-192: (2013).
48. De Silva P.H.G.J, Weerasinghe W.M.C.B, Seresinhe T. Socio-demographic determinants of perceived influences on household food choice in Galle, Kalutara, Matara and Moneragala Districts. *Pakistan Journal of Social Sciences*; **8**(3): 142-145: (2011).
49. Missagia S.V, De Oliveira R.S and De Rezende C.D. Food Choice Motives and Healthy Eating: Assessing Gender differences. *Encontro da ANPAD*; 1-13: (2012).
50. Neumark-Sztainer D, Sherwood N.E, French S.A et al. Weight control behaviors among adult men and women: cause for concern? *Obesity Research*; **7**(2): 179-188: (1999).
51. Kuchler F and Lin B-H. The influence of individual choices and attitudes on adiposity. *International Journal of Obesity*; **26**: 1017-1022: (2002).
52. Paeratakul S, Popkin B.M, Keyou G et al. Changes in diet and physical activity affect the body mass index of Chinese adults. *International Journal of Obesity*; **22**: 424-431: (1998).
53. Hur I, Jang M-J and Oh K. Food and Nutrient Intakes According to Income in Korean Men and Women. *Public Health and Research Perspectives*; **2**(3): 192-197: (2011).
54. Bhurosy T, Jeewon R. Food habits, socioeconomic status and body mass index among premenopausal and post-menopausal women in Mauritius. *Journal of Human Nutrition and Dietetics*; **26**(Suppl. 1):114-

- 122:(2013).
55. Fokeena W.B and Jeewon R. Is There an Association between Socioeconomic Status and Body Mass Index among Adolescents in Mauritius? *The Scientific World Journal*; (2012).
 56. Drewnowski A, Specter S.E. Poverty and obesity: the role of energy density and energy costs. *American Journal of Clinical Nutrition*; **79**: 6-16: (2004).
 57. Chakraborty R, Bose K and Bisai S. Relationship of family income and house type to body mass index and chronic energy deficiency among urban Bengalee male slum dwellers of Kolkata, India. *Journal of Comparative Human Biology*; **60**: 45-57: (2009).
 58. Worsley A. Nutrition knowledge and food consumption: can nutrition knowledge change food behaviour? *Asia Pacific Journal of Clinical Nutrition*; **11**(Suppl): 579-585: (2002).
 59. Edwards A (2011). Food and health concerns in Australia and New Zealand. Available at: <http://www.foodenquirer.com/food-enquirer-channel/articles/food-and-health-concerns-in-australia-and-new-zealand.html>.
 60. Ricciuto L.E and Tarasuk V.S. An examination of income-related disparities in the nutritional quality of food selections among Canadian households from 1986–2001. *Social Science Medicine*; **64**: 186-198: (2007).
 61. Cai L, He J, Song Y et al. Association of obesity with socio-economic factors and obesity-related chronic diseases in rural southwest China. *The Royal Society for Public Health*; **127**: 247-251:(2013).
 62. Xiao Y, Zhao N, Wang H et al. Association between socioeconomic status and obesity in a Chinese adult population. *BMC Public Health*; **355**(13): (2013).
 63. Wilcox S, Sharpe P.A, Turner-McGrievy G et al. Frequency of consumption at fast-food restaurants is associated with dietary intake in overweight and obese women recruited from financially disadvantaged neighborhoods. *Nutrition Research*; **33**: 636-646: (2013).
 64. Pampel F.C, Krueger P.M and Denney J.T. Socioeconomic Disparities in Health Behaviors. *Annual review of sociology*; 349-370: (2010).
 65. Kerver J.M, Yang E.J, Obayashi S et al. Meal and Snack Patterns Are Associated with Dietary Intake of Energy and Nutrients in US Adults. *Journal of the American Dietetic Association*; **106**: 46-53: (2006).
 66. Piernas C and Popkin B.M. Snacking Increased among U.S. Adults between 1977 and 2006. *Nutrition Journal*; **140**: 325-332: (2010).
 67. Forslund H.B, Torgerson J.S, Sjöström L et al. Snacking frequency in relation to energy intake and food choices in obese men and women compared to a reference population. *International Journal of Obesity*; **29**: 711-719: (2005).
 68. Zailiah M.S, Mirnalini K, Safiah M.Y et al. Daily Energy Intake from Meals and Afternoon Snacks: Findings from the Malaysian Adults Nutrition Survey (MANS). *Malaysian Journal of Nutrition*; **14**(1): 41-55: (2008).
 69. Duffey K.J, Pereira R.A and Popkin B.M. Prevalence and energy intake from snacking in Brazil: analysis of the first nationwide individual survey. *European Journal of Clinical Nutrition*; **67**(8): 868-874: (2013).
 70. Hampl J.S, Heaton C.L.B, Taylor C.A. Snacking patterns influence energy and nutrient intakes but not body mass index. *Journal of Human Nutrition and Dietetics*; **16**(1): 3-11: (2003).
 71. Drummond S, Crombie N and Kirk T. A critique of the effects of snacking on body weight status. *European Journal of Clinical Nutrition*; **50**(12): 779-783: (1996).
 73. Viskaal-Van DM, Kok F.J and De Graaf C. Effects of snack consumption for 8 weeks on energy intake and body weight. *International Journal of Obesity*; **34**(2): 319-326: (2010).
 74. Farajian P, Katsagani M and Zampelas A. Short-term effects of a snack including dried prunes on energy intake and satiety in normal-weight individuals. *Eating Behaviors*; **11**(3): 201-203: (2010).
 75. Smith J.M, Ditschun T.L. Controlling satiety: how environmental factors influence food intake. *Trends in Food Science and Technology*; **20**: 271-277: (2009).
 76. Centers For Disease Control And Prevention (2006) Do Increased Portion Sizes Affect How Much We Eat? 2006. Available at: <http://>

- www.cdc.gov/nccddphp/dnpa/nutrition/pdf/portion_size_research.pdf (accessed 25 March 2014).
77. Jeffery R.W, Rydell S, Dunn C.L et al. Effects of portion size on chronic energy intake. *International Journal of Behavioral Nutrition and Physical Activity*; **27**(4): 1-5: (2007).
78. Rolls B.J, Roe L.S and Meengs J.S. Larger Portion Sizes Lead to a Sustained Increase in Energy Intake Over 2 Days. *Journal of the American Dietetic Association*; **106**(4): (2006).
79. Mccrory M.A, Suen V.M.M and Roberts S.B. Biobehavioral Influences on Energy Intake and Adult Weight Gain. *The Journal of Nutrition*; **132**: 3830-3834: (2002).
80. Swinburn B.A, Caterson I, Seidell J.C et al. Diet, nutrition and the prevention of excess weight gain and obesity. *Public Health Nutrition*; **7**(1A): 123-146: (2004).
81. Ello-Martin J.A, Ledikwe J.H, Rolls B.J. The influence of food portion size and energy density on energy intake: implications for weight management. *American Journal of Clinical Nutrition*; **82**(Suppl): 236-241: (2005).
82. Webber C.B, Sobal J, Dollahite J.S. Physical Disabilities and Food Access Among Limited Resource Households. *Disability Studies Quarterly*; **27**(3): (2007).
83. Dressler H, Smith C. Environmental, Personal, and Behavioral Factors Are Related to Body Mass Index in a Group of Multi-Ethnic, Low-Income Women. *Journal of the Academy of Nutrition and Dietetics*; **113**(12): 1662-1668: (2013).
84. Nagata J.M, Vallengia C.R, Barg F.K et al. Body mass index, socio-economic status and socio-behavioral practices among Tz'utujil Maya women. *Economics and Human Biology*; **7**: 96-106: (2009).
85. Jackson M, Walker S, Forrester T et al. Social and dietary determinants of body mass index of adult Jamaicans of African origin. *European Journal of Clinical Nutrition*; **57**: 621-627: (2003).
86. Yannakoulia M, Panagiotakos D, Pitsavos C et al. Eating patterns may mediate the association between marital status, body mass index, and blood cholesterol levels in apparently healthy men and women from the ATTICA study. *Social Science and Medicine*; **66**: 2230-2239: (2008).
87. Marcellini F, Giuli C, Papa R et al. Obesity and Body Mass Index (BMI) in relation to life-style and psycho-social aspects. *Archives of Gerontology and Geriatrics*; **1**: 195-206: (2009).
88. Horwath C.C. Marriage and diet in elderly Australians: results from a large random survey. *Journal of Human Nutrition and Dietetics*; **2**: 185-193: (1989).
89. World Health Organization/Food and Agriculture Organization (2003) Diet, nutrition and the prevention of chronic diseases. Available at: <ftp://ftp.fao.org/docrep/fao/005/ac911e/ac911e00.pdf> (accessed 6 January 2014).
90. Chaput J, Klingenberg L, Rosenkilde M et al. Physical Activity Plays an Important Role in Body Weight Regulation. *Journal of Obesity*; 1-12: (2011).
91. World Heart Federation. Physical activity: vital to global health. Available at: http://www.world-heart-federation.org/fileadmin/user_upload/children/documents/factsheets/Factsheet_Physical_activity.pdf (accessed 13 March 2014).
92. Melzer K, Kayser B, Saris W.H.M et al. Effects of physical activity on food intake. *Clinical Nutrition*; **24**: 885-895: (2005).
93. Schneider K.L, Spring B and Pagoto S.L. Exercise and energy intake in overweight, sedentary individuals. *Eating Behaviors*; **10**: 29-35: (2009).
94. Westerterp K.R. Physical activity, food intake, and body weight regulation: insights from doubly labeled water studies. *Nutrition Reviews*; **68**(3): 148-154: (2010).
95. Chu A.H.Y, Moy F.M. Association between physical activity and metabolic syndrome among Malay adults in a developing country. *Malaysian Journal of Science and Medicine in Sport*; **17**: 195-200: (2014).
96. Bauman A, Bull F, Chey T et al. The International Prevalence Study on Physical Activity: results from 20 countries. *International Journal of Behavioral Nutrition and Physical Activity*; **21**(6): 1-11: (2009).
97. Kaur G, Bains K and Kaur H. Body Composition,

- Dietary Intake and Physical Activity Level of Sedentary Adult Indian Women. *Food and Nutrition Sciences*; **3**: 1577-1585: (2012).
98. Uthman O and Aremu O. Comparison of physical activity level between overweight/ obese and normal weight individuals: A systematic review. *Internet Journal of Nutrition and Wellness*; **5**(1): (2007).
 99. Thorp A.A, Owen N, Neuhaus M et al. Sedentary Behaviors and Subsequent Health Outcomes in Adults A Systematic Review of Longitudinal Studies, 1996–2011. *American Journal of Preventive Medicine*; **41**(2): 207-215: (2011).
 100. Donnelly J.E, Herrmann S.D, Lambourne K et al. Does Increased Exercise or Physical Activity Alter Ad-Libitum Daily Energy Intake or Macronutrient Composition in Healthy Adults? A Systematic Review. *PLoS ONE*; **9**(1): (2014).
 101. Lim K-C, Kayser-Jones J.S, Waters C et al. Aging, Health, and Physical Activity in Korean Americans. *Geriatric Nursing*; **28**(2): 112-119: (2007).
 102. Hiza H.A.B, Casavale O.K, Guenther P.M et al. Diet Quality of Americans Differs by Age, Sex, Race/ Ethnicity, Income, and Education Level. *Journal of the Academy of Nutrition and Dietetics*; **113**: 297-306: (2013).
 103. Food and Agriculture Organization/ World Health Organization/United Nations University (2001) Human Energy Requirements. Available at: <ftp://ftp.fao.org/docrep/fao/007/y5686e/y5686e00.pdf> (accessed 26 March 2014).
 104. Vlismas K, Stavrinou V and Panagiotakos D.B. Socio-economic status, dietary habits and health-related outcomes in various parts of the world: a review. *Central European Journal of Public Health*; **17**(2): 55-63:(2009).
 105. De Vriendt T , Matthys C, Verbeke W et al. Determinants of nutrition knowledge in young and middle-aged Belgian women and the association with their dietary behavior. *Appetite*; **52**(3): 788-792: (2009).
 106. Roberts K and Marvin K (2011) Knowledge and attitudes towards healthy eating and physical activity: what the data tell us. Oxford: National Obesity Observatory.
 107. Drewnowski A, Monsivais P, Maillot M et al. Low-Energy-Density Diets Are Associated with Higher Diet Quality and Higher Diet Costs in French Adults. *Journal of the American Dietetic Association*; **107**: 1028-1032: (2007).
 108. Kim H.J, Lee K.S, Eom J-S et al. Relation between nutritional risk and metabolic syndrome in the elderly. *Archives of Gerontology and Geriatrics*; **52**: 19-22: (2011).
 109. Rathee M and Hooda A. Nutritional Status In Denture Wearers: A Review. *Internet Journal of Nutrition and Wellness*; **10**(2).
 110. Risonar M.G.D, Rayco-Solon P, Ribaya-Mercado J.D et al. Physical activity, energy requirements, and adequacy of dietary intakes of older persons in a rural Filipino community. *Nutrition Journal*; **8**(19): 1-9: (2009).
 111. Seman K and Abdul Manaf H.I.A.R. Dental Caries experience of elderly people living in “Pondok” in Kelantan. *Archives of Orofacial Sciences*; **2**: 20-25: (2007).
 112. Appollonio I, Carabellese C, Frattola A et al. Influence of dental status on dietary intake and survival in community-dwelling elderly subjects. *Age Ageing*; **26**: 445-455: (1997).
 113. Saarela R.T.K, Lindroos E, Soini H, Hiltunen K, Muurinen S, Suominen M.H. and Pitkälä K.H. Dentition, nutritional status and adequacy of dietary intake among older residents in assisted living facilities. *Gerodontology*, **33**(2) : (2014).
 114. Kossioni A and Bellou O. Eating habits in older people in Greece: The role of age, dental status and chewing difficulties. *Archives of Gerontology and Geriatrics*; **52**: 197-201: (2011).