



Utilization of Dairy Industry Waste-Whey in Formulation of Papaya RTS Beverage

**ANIL PANGHAL¹, VIKAS KUMAR¹, SANJU B. DHULL²,
YOGESH GAT¹ and NAVNIDHI CHHIKARA^{*1}**

¹Department of Food Technology & Nutrition, Lovely Professional University, Jalandhar, India.

²Department of Food Technology, Chaudhary Devi Lal University, Sirsa, India.

Abstract

Fruits and vegetable are highly perishable and are subjected to rapid post harvest losses. Their value addition can enhance shelf life, new product development and finally commodity value in market. The present work was planned to prepare papaya based beverage with whey incorporation (0, 25, 50 and 100%), which is a major environmental pollutant from dairy industry. RTS was evaluated for its nutritional composition, physicochemical attributes like titratable acidity, total soluble solids, total and reducing sugars and sensory quality. Microbial quality was also observed for 60 days. Results revealed that on increasing whey incorporation, nutritional quality was enhanced, but poor sensory and microbial quality was observed. Beverage with 25% addition was found most acceptable with overall acceptability of 8.59 ± 0.21 . Selected beverage has 15.05 ± 0.18 °Brix TSS, $0.30 \pm 0.11\%$ acidity, $5.37 \pm 0.01\text{g}/100\text{g}$ reducing sugar, 14.06 ± 0.46 g/100g total sugar and 5.60 ± 0.02 mg/100g ascorbic acid, thus depicting enhanced nutritional value and this creates a scope for better health, enhanced papaya postharvest value and whey utilization.



Article History

Received: 04 July 2017

Accepted: 25 July 2017

Keywords:

beverage,
whey,
papaya,
waste utilisation.


Introduction

Fruits and vegetables embrace an important status among the health foods as they provide significant amount of nutrients, especially vitamins, minerals, fiber sugars and antioxidants as well as it's refreshing and thirst quenching behavior¹. However, due to highly perishable nature and short shelf life of fruits and vegetables, immediate processing into

preserved products becomes essential to avoid post-harvest losses. India contributes 42% of world production from 30% of the global area under papaya cultivation and is largest producer according to FAO report (2012). Papaya accounts for 6.6% of total crop and covers only 1.9% of the total area under fruit cultivation in India². So Papaya was selected for preparation of fruit drink due to its reasonable price,

CONTACT NAVNIDHI CHHIKARA ✉ navnidhi24@gmail.com 📍 Department of Food Technology & Nutrition, Lovely Professional University, Jalandhar, India.

© 2017 The Author(s). Published by Enviro Research Publishers

This is an  Open Access article licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (<https://creativecommons.org/licenses/by-nc-sa/4.0/>), which permits unrestricted NonCommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

To link to this article: <http://dx.doi.org/10.12944/CRNFSJ.5.2.14>

easy availability and high nutritive value. Papaya is also known as common man's fruit and belongs to Caricaceae family. Papaya possess attractive colour, luscious taste, rich source of minerals like potassium and magnesium, nutrients such as carotenoids, vitamins C, E & flavonoids which acts as antioxidants; vitamins B, folate, pantothenic acid; and fiber³. Nutritionally, almost fruits/fruit drinks are devoid of protein, which is an important for body build, repair and immunity.

Whey is a watery liquid by-product of dairy industry during the preparation of channa, paneer, cheese and casein. World whey production is 180 to 190×106 tonnes/year with an annual increment of 1-2% and only 50% of whey is utilized/processed⁴. Whey contains 45-50% total milk solids, 70% milk sugar (lactose), 20% milk proteins and 70-90% milk minerals and almost all the water soluble vitamins originally present in milk⁵. So, Whey disposal becomes a serious environmental pollutant being loaded with high amount of organic matter. Whey possesses preventive and curative elements responsible for treatment of ailments such as arthritis, anemia and liver complaints⁶. Fruit and dairy/dairy waste based products are attaining considerable attention due to delicious taste, increased energy value and high nutritional value and market for such food products has incredible potential⁷. Whey based

fruits beverages are more suitable for health as compared to other drinks⁸. Production of nourishing pleasant whey based fruit RTS beverages is one of the most promising trends in the utilization of dairy waste, whey. The present study was planned to prepare papaya RTS by incorporating different level of whey and study its storage behavior.

Material and method

Sample Procurement and Preparation

Toned milk and papaya were procured fresh from local market. The whey was obtained using toned milk by simple acid coagulation methods. The hot milk in stainless steel vessel was acidified by simple acid (2% citric acid) coagulation method. Mature and fully ripened papaya fruits were selected. Pulping of papaya was done with method of Lal *et al.*,⁹ and stored at $4 \pm 1^\circ\text{C}$ until use. The RTS beverage was prepared by using whey, water, pulp, sugar and citric acid as per FPO specification¹⁰. Different variants of beverages were formulated using three different levels of whey i.e. 25%, 50%, and 75% by replacing water. The prepared RTS was packed in sterilized bottle (capacity 200 ml) leaving $\frac{1}{2}$ " head space and capped air tight. The steps involved in the preparation of whey based papaya RTS are given in Figure 1. After in-bottle pasteurization




Fig. 1. Flow chart for Whey based Papaya RTS beverage preparation

(80 °C for 10 min) and cooling, RTS was stored at refrigeration temperature for 60 days.

Physicochemical analysis

TSS (using Abbe's hand refractometer), titratable acidity, total and reducing sugars, ascorbic acid¹¹ and microbiological load (Total plate count) of RTS was determined periodically (15 days) over the period of 3 months.

Sensory Evaluation

The fresh and stored beverage samples were organoleptically evaluated by 10 semi-trained members from Department of Food Technology and Nutrition, Lovely Professional University, Jalandhar, India, using Hedonic scale ranging from 1-9 for lowest to highest for color, flavor, taste, and overall acceptability¹².

Statistical analysis

Data obtained from the physico-chemical, microbiological and sensory attributes of papaya whey RTS were analyzed by using Graph Pad Prism

(La Jolla, CA, USA) (version 5.01) software. Results are expressed as means \pm SD. Differences between the means were tested for statistical significance using a 2-way ANOVA and followed by Bonferroni post-hoc test. The significance level was set at 5% ($P < 0.05$) for all calculations¹³.

Results and Discussion

Effect of storage on Physico-chemical characteristics of Whey based Papaya RTS beverage

The changes noted in the physicochemical constituents are given in Table 1.

Total soluble solids

With complete replacement of water with whey, TSS was found to increase from 15.20 \pm 0.35 to 18.15 \pm 0.18. This is due to water replacement with whey as whey contains water, milk solids, lactose, proteins and minerals. On storage, TSS of beverage was enhanced during storage period for all samples (Table 1). However, retention or slight increase in TSS content of juice is desirable for preservation of good

Table 1: Effect of Storage on Physico-chemical characteristics of Whey based Papaya RTS Beverage

Level (%) of water replacement with whey	Storage period (days)	TSS (° Brix)	Acidity (%)	Reducing Sugar (g %)	Total Sugar (g %)	Ascorbic acid (mg %)
Control (0 % whey)	0	14.98 \pm 0.24	0.30 \pm 0.3	5.34 \pm 0.01	13.80 \pm 0.38	5.60 \pm 0.04
	15	15.09 \pm 0.16	0.30 \pm 0.2	5.36 \pm 0.01	13.98 \pm 0.45	5.58 \pm 0.03
	30	15.12 \pm 0.18	0.32 \pm 0.15	5.36 \pm 0.04	14.06 \pm 0.51	5.54 \pm 0.03
	45	15.18 \pm 0.22	0.32 \pm 0.16	5.38 \pm 0.02	14.14 \pm 0.48	5.52 \pm 0.04
	60	15.20 \pm 0.35	0.34 \pm 0.18	5.40 \pm 0.01	14.23 \pm 0.55	5.51 \pm 0.02
A (25% whey)	0	15.05 \pm 0.18	0.30 \pm 0.11	5.37 \pm 0.01	14.06 \pm 0.46	5.60 \pm 0.02
	15	15.22 \pm 0.09	0.32 \pm 0.09	5.37 \pm 0.02	14.14 \pm 0.39	5.59 \pm 0.02
	30	15.65 \pm 0.21	0.32 \pm 0.13	5.37 \pm 0.02	14.22 \pm 0.43	5.57 \pm 0.05
	45	15.79 \pm 0.17	0.33 \pm 0.12	5.39 \pm 0.03	14.30 \pm 0.51	5.56 \pm 0.09
	60	15.85 \pm 0.19	0.33 \pm 0.13	5.42 \pm 0.03	14.38 \pm 0.42	5.54 \pm 0.03
B (50% whey)	0	15.88 \pm 0.13	0.30 \pm 0.17	5.40 \pm 0.03	14.09 \pm 0.13	5.60 \pm 0.05
	15	16.05 \pm 0.17	0.32 \pm 0.11	5.42 \pm 0.02	14.19 \pm 0.26	5.60 \pm 0.01
	30	16.38 \pm 0.15	0.34 \pm 0.08	5.45 \pm 0.05	14.22 \pm 0.19	5.58 \pm 0.06
	45	16.95 \pm 0.19	0.34 \pm 0.12	5.46 \pm 0.02	14.37 \pm 0.24	5.55 \pm 0.05
	60	17.25 \pm 0.21	0.36 \pm 0.13	5.48 \pm 0.03	14.49 \pm 0.44	5.52 \pm 0.02
C(100% whey)	0	17.03 \pm 0.13	0.30 \pm 0.12	5.45 \pm 0.05	14.30 \pm 0.22	5.60 \pm 0.04
	15	17.25 \pm 0.16	0.32 \pm 0.16	5.46 \pm 0.04	14.48 \pm 0.32	5.58 \pm 0.03
	30	17.78 \pm 0.08	0.34 \pm 0.19	5.46 \pm 0.03	14.62 \pm 0.17	5.57 \pm 0.03
	45	18.02 \pm 0.12	0.35 \pm 0.12	5.49 \pm 0.03	14.78 \pm 0.24	5.55 \pm 0.04
	60	18.15 \pm 0.18	0.37 \pm 0.13	5.52 \pm 0.02	14.93 \pm 0.18	5.53 \pm 0.02

juice quality during storage¹⁴. This small increment in TSS might be due to polysaccharides hydrolysis into monosaccharide and soluble disaccharides¹⁵.

Acidity

A gradual increase in the acidity of RTS beverage was observed during storage. The average initial acidity of papaya RTS was 0.30 ± 0.3 and increased to $0.34 \pm 0.18\%$. Similar results has been reported by Saravana *et al.*,¹⁶ in whey based jack fruit RTS;

Panghal *et al.*,¹⁷ in whey based strawberry RTS. During storage, degradation of sugar into carboxyl acids, lactose into lactic acid or pectin into pectinic acid is responsible for slight increase in titratable acidity¹⁸. Whey protein might get converted into amino acids which accounts for small increase in titratable acidity¹⁹.

Total and reducing sugars

Sugars were present in form of fruit sugar, lactose




Fig. 2: Total plate count of Whey based Papaya RTS Beverage during storage

Table 2 Effect of Storage on Organoleptic Quality of Whey based Papaya RTS Beverage

Level (%) of water replacement with whey	Storage period (days)	Color	Flavor	Taste	Overall acceptability
Control (0 % whey)	0	9.00±0.17	9.00±0.21	8.99±0.13	9.00± 0.43
	15	8.82±0.11	8.85±0.19	8.78±0.25	8.82±0.36
	30	8.69±0.19	8.68±0.08	8.06±0.33	8.48±0.29
	45	8.53±0.24	8.51±0.31	7.54±0.16	8.19±0.22
	60	8.29±0.13	8.26±0.25	7.06±0.22	7.87±0.39
A (25% whey)	0	8.13±0.09	8.83±0.18	8.82±0.18	8.59±0.21
	15	7.98±0.11	8.45±0.24	8.58± 0.37	8.34±0.19
	30	7.59±0.13	8.16±0.16	8.12±0.22	7.96±0.24
	45	7.26±0.26	7.92±0.11	7.89±0.29	7.69±0.33
	60	7.00± 0.34	7.13±0.21	7.45±0.13	7.19±0.18
B (50% whey)	0	7.92±0.27	8.12±0.25	8.68±0.17	8.24±0.13
	15	7.02±0.15	7.59±0.18	8.08±0.23	7.56±0.16
	30	6.78±0.13	7.16±0.22	7.68±0.28	7.21±0.41
	45	6.39±0.08	6.89± 0.53	7.05±0.31	6.78±0.13
	60	6.05±0.18	6.43±0.23	6.49±0.18	6.32±0.39
C(100% whey)	0	7.16±0.13	7.76±0.19	8.18±0.27	7.65± 0.43
	15	6.85±0.24	7.15±0.36	7.53±0.34	7.18±0.31
	30	6.54±0.17	6.83±0.13	7.17±0.21	6.85±0.24
	45	6.13±0.24	6.54± 0.28	6.42±0.18	6.36±0.13
	60	5.82±0.35	6.03±0.17	6.01±0.38	5.95±0.19

and added sugar in all formulations. So with increase in level of whey incorporation from 0 to 75%, total (13.80±0.38, 14.06±0.46, 14.09±0.13, 14.30±0.22) and reducing sugars (5.34±0.01, 5.37±0.01, 5.40±0.03, 5.45±0.05) were increased (Table 1). The reducing sugars and total sugars increased during storage of 60 days at refrigeration temperature for all the treatments, this might be due to hydrolysis of polysaccharides like starch, cellulose, pectin, etc. and conversion into simple sugars (glucose, fructose). However, rise in total sugars was non-significant and similar results have been reported by²⁰, Krishnaveni *et al.*,²¹ in Jack fruit RTS, Kausar *et al.*,²² in cucumber–melon functional drink, Majumdar *et al.*,²³ in bottled gourd–basil leave juice.

Ascorbic Acid

Ascorbic acid also did not show any significant degradation after storage at refrigeration temperature (Table 1). Ascorbic acid content decreases with the progression of storage time which might be due to oxidation reaction by residual oxygen, followed by decomposition which might have been accelerated due to the storage time. These results are in well coherence with Saravana *et al.*¹⁶; Ibrahim ²⁴.

Effect of storage on Microbial quality of Whey based Papaya RTS beverage

The microbial load of RTS was analyzed periodically during storage (Figure 2) and it showed that there was negligible increase in microbial load and beverage was safe for consumption. Total plate count was comparatively higher as level of whey addition increased due to more availability of nutrients in

whey (Figure 2). Whey contains lactose sugar and proteins favorable for microorganism growth. Similar results have been reported by Sakhale *et al.*,²⁵ for whey-based RTS from mango.

Effect of storage on Organoleptic Evaluation of RTS beverage

Beverage formulated with different whey combinations (0, 25, 50, 100%) were evaluated at regular intervals for various sensorial attributes by using 9 point hedonic scale. The sensory scores for various parameters (color, flavor, taste and overall acceptability) are presented in Table 2.

On increasing whey level, all sensory parameters were decreased. Color values decreased from 9.00±0.17 to 7.16±0.13. Flavor and taste scores were also declined to 7.76±0.19 and 8.18±0.27 respectively. Thus, overall acceptability decreased from 9.00± 0.43 to 7.65± 0.43 with increase of whey level from 0 to 100%. Addition of whey is responsible for dull whitish appearance of beverage, reduced flavor, taste and thus overall acceptability. However, beverage with 25% whey was quite comparable to control sample and was found acceptable as compared to other whey incorporated beverages (Figure 3).

During storage, overall acceptability declined for all treatments, however, quality degradation was higher in case of high whey level. Whey accounts for more microbial growth, reduced acidity and off flavor. So, during storage there is immense loss of color due to browning, increase in acidity changes taste and flavor of beverage. Thus overall acceptability of the samples decreased during storage (Table 2). Similar




Fig. 3: Overall acceptability of Whey based Papaya RTS Beverage during storage

results have been reported by Sakhale *et al.*,²⁵ for whey-based RTS from mango.

Conclusion

The study revealed that whey can be a suitable ingredient for development of whey based healthy beverages. However, addition of whey was found acceptable only up to 25% in terms of nutritional information and consumer acceptability. Distinct papaya flavor is capable to mask unpleasant taste

and flavor of whey. This approach can be suitable for proper utilization of dairy waste whey and beneficial for papaya value addition.

Acknowledgement

This research was supported by Lovely Professional University, Punjab. We thank our colleagues from Lovely Professional University and Chaudhary Devi Lal University, Haryana who provided insight and expertise that greatly assisted the research.

Reference

1. Yadav DN, Vishwakarma RK, Borad S, Bansal S, Jaiswal AK, Sharma M. Development of protein fortified mango based ready-to-serve beverage. *Journal of Food Science and Technology*; 2016. **53**(10):3844-52.
2. Sharma SK, Mitra SK, Saran S. Papaya production in India-history, present status and future prospects. In XXIX International Horticultural Congress on Horticulture: Sustaining Lives, Livelihoods and Landscapes (IHC2014): IV 1111 2014 Aug 17 (pp. 87-94).
3. Ramachandran P, Nagarajan S. Quality Characteristics, Nutraceutical Profile, and Storage Stability of Aloe Gel-Papaya Functional Beverage Blend. *International Journal of Food Science*; 2014. doi:10.1155/2014/847013
4. Román A, Vatai G, Ittész A, Kovács Z, Czermak P. Modeling of diafiltration processes for demineralization of acid whey: an empirical approach. *Journal of Food Process Engineering*; 2012. **35**(5):708-14.
5. Horton BS. Whey processing and utilization. *Bulletin IDF* 1995. 308:2-6
6. Cruz AG, Ana ASS, Macchione MM, Teixeira AM, Schmidt FL. Milk drink using whey butter cheese (Queijo manteiga) and Acerola juice as a potential source of vitamin C. *Food and Bioprocess Technology*; 2009. **2**: 368-373.
7. Ismail AE, Abdelgader MO, Azhari Ali A. Microbial and Chemical Evaluation of Whey-Based Mango Beverage. *Advance Journal of Food Science and Technology*; 2011. **3**: 250-253.
8. Kumar R S. Whey beverage, A review. *Beverage and food World*, 2005. 58-60.
9. Lal G, Siddappa GS, Tandon GL. Preservation of fruits and vegetables. Indian council of agriculture research (ICAR), New Delhi. 1967. 294-297.
10. Gridharilal GS, Siddappa, Tandon GL. Preservation of fruits and vegetables. Indian Council of Agricultural Research, New Delhi. 1998.
11. Ranganna, S. Manual analysis of fruit and vegetable products. 2nd Edn. Tata McGraw Hill Publishing Co., Ltd., New Delhi. 1995.
12. Amerine MA, Pangborn RM, Roessler EB. Principles of sensory evaluation of food. New York, London: Academic Press. 1965.
13. Saliganti, Vamshi, Kapila R, Kapila S. Consumption of probiotic *Lactobacillus rhamnosus* (MTCC: 5897) fermented milk plays a key role on newborn mice immune system development during suckling-weaning transition: Probiotics for early immune maturation. *Microbiology and Immunology*, 2015. **114**(8):1168-79.
14. Bhardwaj RL, Pandey S. Juice blends—a way of utilization of under-utilized fruits, vegetables, and spices: a review. *Critical reviews in food science and nutrition*; 2011. **51**(6):563-70.
15. Datey SP, Raut VU. Physico-chemical changes in mango pulp at ambient storage in glass containers. *Green Farming*; 2009. **2**(10):713-714.
16. Saravana R, Kumar, Manimegalai G. Studies on whey based jack fruit RTS beverage. *Beverage and Food World*; 2002. **29**(1): 57-

- 58.
17. Panghal A, Dhull N, Navnidhi, Khatkar BS. Whey Based Strawberry Ready to Serve (RTS) Beverage. *Beverage and Food World*; 2009. **36**(04) (28-30)
18. Kinh SA, Dunne CP, Hoover DG. Preparation and preservation of apple pulp with chemical preservatives and mild heat. *Journal of Food Protection*; 2001. **28**(6): 111-114.
19. Sikder B, Sarkar K, Ray PR, Ghatak PK. Studies on shelf life of whey Based Mango Beverage. *Beverage and Food World*; 2001. **28**: 53-54.
20. Lanjhiyana R, Sharma PK, Shukla N. Studies on effect of chemical preservatives on physico-chemical changes of beverages in lime and ginger juice with their combinations. *Journal of Horticulture Sciences*; 2010. **5**(2): 151-154.
21. Krishnaveni A, Manimegalai G, Saravanakumar R. Storage stability of jack fruit (*Artocarpus heterophyllus*) RTS beverage. *Journal of Food Science and Technology*; 2001. **38**(6):601-2.
22. Kausar H, Saeed S, Mushtaq Ahmad M, Salam A. Studies on the Development and Storage Stability of Cucumber-Melon Functional Drink. *Journal of Agricultural Research*; 2012. 1:**50**(2).
23. Majumdar TK, Wadikar DD, Vasudish CR, Premavalli KS, Bawa AS. Effect of storage on physico-chemical, microbiological and sensory quality of bottlegourd-basil leaves juice. *American Journal of Food Technology*; 2011. **6**(3):226-34.
24. Ibrahim, M. A. Effect of different storage condition on pH and vitamin C content in some selected fruit juices (pineapple, pawpaw and watermelon). *International Journal of Biochemistry Research and Review*; 2016. **11**(2): 1-5.
25. Sakhale BK, Pawar VN, Ranveer RC Studies on the Development and Storage of Whey based RTS Beverage from Mango cv. Kesar. *Journal of Food Processing and Technology*; 2012. **3**:148.